

# Antimalarial plants of northeast India: An overview

Rama Shankar, Sourabh Deb<sup>1</sup>, B. K. Sharma<sup>2</sup>

Ayurveda Regional Research Institute, Itanagar, <sup>1</sup>Regional Centre, NAEB, NEHU, Shillong, <sup>2</sup>North Eastern Institute of Folk Medicine, Department of AYUSH, Pasighat, Arunachal Pradesh, India

## ABSTRACT

The need for an alternative drug for malaria initiated intensive efforts for developing new antimalarials from indigenous plants. The information from different tribal communities of northeast India along with research papers, including books, journals and documents of different universities and institutes of northeast India was collected for information on botanical therapies and plant species used for malaria. Sixty-eight plant species belonging to 33 families are used by the people of northeast India for the treatment of malaria. Six plant species, namely, *Alstonia scholaris*, *Coptis teeta*, *Crotolaria occulta*, *Ocimum sanctum*, *Polygala persicariaefolia*, *Vitex peduncularis*, have been reported by more than one worker from different parts of northeast India. The species reported to be used for the treatment of malaria were either found around the vicinity of their habitation or in the forest area of northeast India. The most frequently used plant parts were leaves (33%), roots (31%), and bark and whole plant (12%). The present study has compiled and enlisted the antimalarial plants of northeast India, which would help future workers to find out the suitable antimalarial plants by thorough study.

**Key words:** Alkaloids, malaria, medicinal plants, mosquito repellents, northeast India, traditional knowledge of medicine

## INTRODUCTION

Malaria is caused by single-celled protozoan parasites called *Plasmodium* and transmitted to man through the *Anopheles* mosquito. It is one of the major fatal diseases in the world, especially in the tropics, and is endemic in some 102 countries, with more than half of the world population at risk with fatality rates being extremely high among young children below 5 years of age.<sup>[1]</sup> The World Health Organization estimates that there are between 300 and 500 million new cases of malaria worldwide, every year, mostly in Africa, Asia, South Pacific Islands and South America, which causes at least 1 million deaths annually. In spite of control programs in many countries, there has

been very little improvement in the control of malaria, and infections can reduce the effectiveness of labor and can lead to both economic and human losses. Control of malaria is complex because of the appearance of drug-resistant strains of *Plasmodium* and with the discovery that man becomes infested with species of simian (monkey) malaria.<sup>[1]</sup> At the same time, the *Anopheles* mosquitoes have developed resistance to many insecticides.<sup>[2]</sup>

Spread of multidrug-resistant strains of *Plasmodium* and the adverse side effects of the existing antimalarial drugs have necessitated the search for novel, well-tolerated and more efficient antimalarial drugs that kill either the vector or the parasite. The use of plant-derived drugs for the treatment of malaria has a long and successful tradition. The first antimalarial drug was quinine, isolated from the bark of *Cinchona* species (Rubiaceae) in 1820. It is one of the oldest and most important antimalarial drugs, which is still used today. In 1940, another antimalarial drug, chloroquine, was synthesized and is being used for the treatment of malaria.<sup>[3]</sup> Unfortunately, after an early success, the malarial parasite, especially *Plasmodium falciparum*, also became resistant to chloroquine. Treatment of chloroquine-resistant malaria was done with alternative drugs or drug combinations, which were rather expensive and sometimes toxic.<sup>[4]</sup> The extract of the bark and leaves of *Azadirachta indica* has also been used in Thailand and Nigeria as an antimalarial for a long time. Charaka in 300 BC and Susruta in 200 BC reported the antimalarial and antipyretic activity of this

### Address for correspondence:

Dr. Rama Shankar,  
Ayurveda Regional Research Institute, Itanagar – 791 111,  
Arunachal Pradesh, India. E-mail: rshankar58@gmail.com

Received: 10-Jan-2011

Revised: 22-Mar-2011

Accepted: 29-Mar-2011

### Access this article online

#### Quick Response Code:



#### Website:

[www.jaim.in](http://www.jaim.in)

#### DOI:

10.4103/0975-9476.93940

species. Hence, it is clear that the main drugs developed for malaria and used until now (quina alkaloid derived drugs and artemisinin) were discovered based on traditional use and ethnomedical data.

In Ghana, several plant species including *Az. indica* A. Juss, (Meliaceae), *Cryptolepis sanguinolenta* (Lindl.) Schtr. (Asclepidaceae), *Nauclea latifolia* Sm. (Rubiaceae), *Ocimum viride* Willd. (Lamiaceae), *Alstoni boonei* De Willd (Apocynaceae), *Morinda lucida* Benth. (Rubiaceae), *Nyctanthes arbor-tristis* L. (Nyctaginaceae) and *Timospora cordifolia* (Willd.) Miers ex Hook f. and Thomas<sup>[5,6]</sup> are used in the treatment of malaria. The list of antimalarial plants of India has not yet been completely searched out and it is an urgent need to compile this data. The aim of this study was to compile the antimalarial plants reported by different workers from northeast India. The present database of antimalarial plants would help the future phytochemist to evaluate the best antimalarial plants and it would be possible to formulate the most effective medicine from this region of the world. It might, therefore, be useful to test the antibacterial, antiviral and anti-inflammatory activities of these groups of plants. The present survey has provided information about the range of species of plants used in the treatment of malaria in northeast India. Accordingly, researchers should consider the ethnomedical information of all species before deciding which kind of screening should be used in the search for an antimalarial.

## MATERIALS AND METHODS

All primary ethnobotanical studies from books and journals, research papers of different universities and institutes of northeast India were collected for information about botanical therapies and plant species used for malaria. Local traditional healers were interacted with for confirmation and validation as far as possible. Any data or references to plants used for malaria were carefully inserted into a template and botanical name and classification were re-examined and confirmed with the flora of northeast India and flora of India. For compliance of study, authors had interaction with traditional healers through interviews in the prominent communities of Adi (4), Apatani (2), Khampti (3), Mishmis (4), Nyishi (4), Monpa (6), Nocte (2), and Sherdukpens (2) in Arunachal Pradesh, Mizo (4) in Mizoram, Aos (3) in Nagaland, and Nepalis and Lepcha (2) in Sikkim. As per the available literature, various folklores treating malaria in different communities have been reported.<sup>[7-27]</sup>

## RESULTS AND DISCUSSION

After thorough literature survey using the above method, it

can be confirmed that 68 species of plants belonging to 33 families are used by the people of northeast India for the treatment of malaria [Table 1]. Of the 33 families studied, Verbenaceae (5), Acanthaceae (3), Asteraceae (5), Rubiaceae (3), Rutaceae (4), Lamiaceae (5), and Euphorbiaceae (3) are predominant in terms of the number of species used to treat malaria. Six plant species, namely, *Alstonia scholaris*, *Coptis teeta*, *Crotalaria occulta*, *Ocimum sanctum*, *Polygala persicariaefolia*, and *Vitex peduncularis*, have been reported by more than one worker from different parts of northeast India. The species reported to be used for the treatment of malaria were either found around the vicinity of their habitation and in the forest area of northeast India. More than 20 authors have reported antimalarial plants and the author himself has reported 10 antimalarial plants from different parts of Arunachal Pradesh.<sup>[7-9]</sup> Similarly 9, 12, 4 and 2 species from Assam, Manipur, Mizoram and Sikkim respectively.<sup>[10-12,15-23]</sup>

Most of the plants were reported from Assam, Arunachal Pradesh, Manipur and Mizoram, whereas Nagaland and Tripura have been less explored by any author. The plants recorded in this survey were used traditionally for the treatment of malaria and its symptoms. Majority of the plants were used as decoctions and some plants were used both internally and externally. Herbs and shrubs were found to be predominantly used as antimalarial drugs, and the most frequently used plant parts were leaves (33%), roots (31%), and bark and whole plant (12%). The enormous frequency of the usage of leaves in traditional preparations is related to their abundant availability and easy collection.

Information from traditional healers of Assam Ayurveda Regional Research Institute, Itanagar, revealed that they had used pills of *Kalmegh* (*Andrographis paniculata*), stem bark of *Latakaranja* (*Caesalpinia crista*) and *Guduchi* (*T. cordifolia*). Some species like *Holoptelea integrifolia* Planch, *T. cordifolia* (Willd.), *Calotropis procera* (Ait.) R. Br., *Nerium indicum* Mill., *Ajuga bracteosa* Wall., *Leucas cephalotes* Spreng., *Enicostemma byssopifolium* (Willd.) Verdoorn, *Vernonia cinerea* Less., *Justicia adhatoda* Linn., *Orthosiphon pallidus* Royle ex Benth., *Pongamia pinnata* (L.) Merr., *Nyctanthes arbor-tristis* L., *Calotropis gigantea* (L.) R. Br., *Capsicum annum* L., *Phyllanthus fraternus* L., *Plectranthus* sp., *Elephantopus scaber* Linn., *Combretum decandrum* Roxb., *Holarrhena antidysenterica* Wall., *Cleome viscosa* L., *Vernonia roxburghii* Less., *Achyranthes aspera* L. are also available in northeast India, but any report on their use in any part of northeast India has not yet been published.<sup>[28-30]</sup>

The knowledge of plants used in the treatment of malaria in northeast India, combined with the high level of correlation found with the uses of these plants (or related species) in diverse parts of India, indicates the inheritance of our ancestral knowledge throughout the country. It represents

**Table 1: List of antimalarial plants reported from northeast India**

Name of the plant	Family	Vern. name	Parts used	Methodology	References
<i>Acacia farnesiana</i> (L.) Willd	Mimosaceae	<i>Tarua kadam</i> (Ass)	Bark		[10]
<i>Acorus calamus</i> L.	Araceae	<i>Bach</i> (Beng), Sweet flag (Eng)	Rhizome	If taken with quinine, stops remittent fever	[15]
<i>Adhatoda zeylanica</i> Medicus	Acanthaceae	<i>Kawldai</i> (Mi)	Leaf	The leaves are boiled and the water is used for bathing and the leaf paste is applied on the whole body as an effective cure for chronic fever/malaria	[12]
<i>Alstonia scholaris</i> R.Br.	Apocynaceae	<i>Tun tong</i> (Khamti), <i>Chatiana</i> (Assamese), <i>Thamrita</i> (Mi)	Bark	Bark infusion is given once a day	[16], [17], [18], [7], [10], [19]
<i>Andrographis paniculata</i> Wall. Ex Nees	Acanthaceae	<i>Gokur</i> (Beng), <i>Kalmegh</i> (S), <i>Hnakapui</i> (Mi), <i>Vubati</i> (Man)	Leaf	Crushed raw leaves are taken orally for 2 days twice with half glass of milk	[20]
<i>Artemisia nilagirica</i> (C.B. Clarke) Pamp.	Asteraceae	<i>Koken</i> (Nyishi), <i>Sai</i> (Mi), <i>Laibakngou</i> (Man), <i>Nagdona</i> , <i>Tongloti</i> (Ass)	Leaf	Decoction of leaves is given	[21]
<i>Asplenium adiantoides</i> C. Chr.	Aspleniaceae	<i>Ruimangma</i> (Man)	Plant		[22]
<i>Aster amellus</i> L.	Asteraceae		Root		[22]
<i>Berberis aristata</i> DC.	Berberidaceae	<i>Daru Haridra</i> (S), <i>Drauhaldi</i> (Beng)	Root	The root bark is used as tonic	[15]
<i>Betula alnoides</i> Buch.-Ham	Betulaceae	<i>Hriang</i> (Mi), <i>Bhujpattra</i> (Hi)	Bark	Decoction is taken	[17]
<i>Brucea javanica</i> (Linn.) Merr.	Simaroubaceae	<i>Heining</i> (Man), <i>Tammu</i> (Rongmei)	Fruit		[11]
<i>Carica papaya</i> L.	Caricaceae	<i>Papeya</i> (Beng)	Leaf		[10]
<i>Cinchona officinalis</i> Linn f.	Rubiaceae		Bark	The bark of the tree is grounded into powder and then it is boiled in water and fed to the patient	[23], [19]
<i>Cinnamomum bejolghota</i> (Buch.-Ham)	Lauraceae	<i>Tezpta</i> (Mi)	Bark and leaf	The bark and leaves are boiled with the leaves of <i>Anacolosia crassipes</i> . The water is used for bathing, the steam inhaled and the water taken internally	[12]
<i>Cissampelos pareira</i> L.	Menispermaceae	<i>Tubuki lot</i> (As), <i>Papurilota</i>	Root	Juice is used	[10]
<i>Citrus medica</i> L.	Rutaceae	<i>Baranimbu</i> (Beng)	Fruit	Juice is used	[10]
<i>Citrus sinensis</i> (L.) Osbeck	Rutaceae	<i>Musambi</i> (M, H and B) Sweet orange <i>Serthlum</i> (M), <i>Kamalanimbu</i> (H)	Leaf	Decoction is taken	[24]
<i>Clausena excavata</i> Burm. f.	Rutaceae	<i>Bhant</i> (H)	Leaf	Juice rubbed to alleviate muscular pain	[25]
<i>Clerodendron infortunatum</i> Gaertn.	Verbenaceae	Assam	Root and leaf		[26]
<i>Clerodendrum colebrookoianum</i> Walp.	Verbenaceae	<i>Nephaphu</i> (Ass), Ar	Leaf	Decoction is given to cure	[25]
<i>Clerodendrum serratum</i> (L.) Moon	Verbenaceae	<i>Barangi</i> (H)	Root		[10]
<i>Coptis teeta</i> Wall	Ranunculaceae	<i>Mishmi teeta</i>	Root, rhizome	It is administered orally at a dose of 150 g thrice a day	[27], [25], [7]
<i>Crotolaria occulta</i> Grab	Fabaceae		Plant	Plant juice taken with warm water	[7], [25]

Table 1 (contd...)

Shankar, *et al.*: Antimalarial plants of northeast India

Table 1 (contd...)

Name of the plant	Family	Vern. name	Parts used	Methodology	References
<i>Croton tiglium</i> L.	Euphorbiaceae	Jaiphal (H)	Leaf, flower	Plant powder is consumed with a glass of water twice a day till cured	[20]
<i>Croton caudatus</i> Geisel	Euphorbiaceae		Root		[10]
<i>Cynoglossum glochidion</i> Wall.	Boraginaceae	Nyumli Mento (Monpa)	Root	Root pounded or powdered and mixed with water. 10 g is taken twice daily	[7]
<i>Datura metel</i> L.	Solanaceae	Dhatura (H)	Seed, leaf and root	In fever with catarrhal and cerebral complication	[21]
<i>Dichroa febrifuga</i> Lour.	Saxifragaceae	Khawsik-damdawi (Mi), Basak (H)	Root and leaf	Roots and leafy tops are used in malarial fever. Therapeutic activity is due to quinazoline derivatives	[21]
<i>Elsholtzia blanda</i> Benth	Lamiaceae	Papit Namdung (Adi) Pheiri (Tankhul), Bantulsi (A)	Leaf	It is a mosquito repellent	[25]
<i>Gomphostemma parviflora</i> Wall.	Lamiaceae		Leaf		[10]
<i>Halenia elliptica</i> D. Don	Gentianaceae	Qing ye dan (Mon)	Plant	Taken orally in malarial fever	[7]
<i>Hedyotis scandens</i> Roxb.	Rubiaceae	Bakrelara (Nep) Khangbai taak (Rongmei)	Root and leaf	Infusion of the roots and leaves is taken as an effective remedy	[12]
<i>Helianthus annuus</i> L.	Asteraceae	Numitlei (M), Surajmukhi (B and H)	Leaf and flower	Decoction of plant parts along with honey is prescribed	[11], [10]
<i>Homonoia riparia</i> Lour.	Euphorbiaceae	Tuipui sulhla (A)	Wood	Wood infusion is given	[11]
<i>Hydrangea macrophylla</i> (Thunb.) Ser.	Saxifragaceae		Leaf, root	Said to be more potent than quinine	[11]
<i>Impatiens angustifolia</i> Blume	Balsaminaceae		Leaf	Paste is given	[25]
<i>Lantana camara</i> L.	Verbenaceae	Hlingpangpar (Mi)	Plant	Plant decoction is given	[12]
<i>Magnolia grandiflora</i> L.	Magnoliaceae	U-thambal (M), Andachampa (H and B)	Bark		[11]
<i>Melodinus monogynus</i> Roxb.	Apocynaceae		Leaf, wood and root	Contains a narcotic poison. Used as antimalarial drug	[21]
<i>Mesona wallichiana</i> Benth.	Lamiaceae		Root	Boiled extract is given	[28]
<i>Nasturtium officinale</i> Br.	Brassicaceae	Chhu-ra (Nyishi)	Plant	Plant decoction (2–3 teaspoonful) taken twice daily	[7]
<i>Ocimum sanctum</i> L.	Lamiaceae	Tulasi (H and S)	Root	Decoction is given as diaphoretic	[17], [22]
<i>Ocimum tenuiflorum</i> L.	Lamiaceae		Root		[14]
<i>Passiflora nepalensis</i> Walp.	Pasifloraceae	Nauawimu (M)	Root	Decoction is taken	[24]
<i>Picrasma javanica</i> Bl.	Simaroubaceae	Thingdamdawi/Khwsik-damdawi thing	Bark	The inner bark is very bitter like cinchona. An infusion of the inner coat of bark is taken orally in lieu of quinine at a dose of tablespoonful (10 ml) twice daily	[12]
<i>Piper mullesua</i> Buch. Ham.	Piperaceae	Pippali (Adi)	Leaf, fruit	Dried plant is consumed during malaria and cough	[20]
<i>Polygala persicariaefolia</i> DC.	Polygalaceae		Plant	Boiled and decoction is given	[11], [25]

Table 1 (contd...)

Table 1 (contd...)

Name of the plant	Family	Vern. name	Parts used	Methodology	References
<i>Randia fasciculata</i> (Roxb.) DC.	Rosaceae	<i>Chhawntan</i> (M), <i>Horu-moyan</i> , <i>pulikaint</i> (As)	Leaf	Leaf mixed with Piper nigrum and their boiled juice extract is given	[26]
<i>Rubus ellipticus</i> Sm.	Rosaceae	<i>Hmutau</i>	Leaf	Root decoction used	[18]
<i>Rubus</i>				Leaf mixed with Piper nigrum and their boiled juice extract is given	[18]
<i>Satyrium nepalense</i> D. Don	Orchidaceae		Tuber	Consumed as tonic	[25]
<i>Sida rhombifolia</i> L.	Malvaceae	<i>Boriala</i> (Ass)	Root	Boiled extract is given	[18]
<i>Solanum vairum</i> Cl.	Solanaceae	<i>Thitbyake</i> (Nyishis)	Root	Decoction of root is used	[7]
<i>Solanum torvum</i> Sw.	Solanaceae	<i>Gota begun</i>	Fruit	Burnt fruits are consumed	[20], [10]
<i>Stephania japonica</i> Miers.	Menispermaceae	<i>Rajpatha</i> (H)	Tuber	Sun-dried tuber powder is administered orally with boiled water twice a day for more than 4 days till malaria is cured	[20]
<i>Strobilanthes auriculatus</i> Nees	Acanthaceae	<i>Ramting</i> (M), <i>Rum</i> (Ass)	Leaf	Pounded leaves rubbed on the body during the cold stage of intermittent fever.	[21]
<i>Swertia dilatata</i> Wall.	Gentianaceae	<i>Sirota</i> (Ass)	Root	Powdered root is administered	[25]
<i>Swertia nervosa</i> Wall.	Gentianaceae	<i>Sirota</i> (Ass)	Plant	Decoction of plant is given	[25]
<i>Taraxacum officinalis</i> Wigg.	Asteraceae	<i>Pitachumki</i> (Beng), <i>Dudal</i> (Hindi)	Plant	Powder is used	[25]
<i>Thalictrum foliolosum</i> L.	Ranunculaceae	<i>Pilijari</i> (H), <i>Gurbiani</i> (Beng)	Rhizome and root	Extract is a bitter tonic	[7]
<i>Vandellia sessiliflora</i> Benth.	Scrophulariaceae		Plant	Decoction of whole plant is used	[7]
<i>Vitex peduncularis</i> Wall	Verbenaceae	<i>Thing-khawi-lu</i> (Mizo)	Bark, leaf, stem	The bark is crushed and boiled. The steam vapor is inhaled by a patient suffering from malarial fever; infusion of leaves or of root bark or young stem bark is useful in malaria and black water fever	[12], [17], [25]
<i>Picrorhiza kurroa</i> Benth.	Scrophulariaceae		Root	Pounded in water and given	[15]
<i>Xanthium strumarium</i> L.	Asteraceae	<i>Agara</i> (Ass)	Leaf		[14]
<i>Zanthoxylum hamiltonianum</i> Wall.	Rutaceae		Root and bark		[27]

(Ass) - Assamese, (H)- Hindi, (Beng)- Bengali, (M) - Manipuri, (Mi) - Mizo, (S)- Sanskrit

sometimes the only available alternative malarial treatment in remote communities of the northeast India. Four species from Assam, Arunachal Pradesh, Meghalaya and Mizoram were also found to be used as mosquito repellents [Table 2]. Though species like *Homalomena aromatica*, *Ocimum gratissimum*, *Elsoltzia blanda*, *Eucalyptus globules* are written as repellents, whether these plants are repellents or insecticides or both has not yet been sufficiently proved. Local people of this region used these plants as a substitute for DDT and other insecticides, as it is well known that DDT and other insecticides have adverse effects on environment and human health. Several classes of the secondary plant metabolites are responsible for antimalarial activity, but the most important and diverse biopotency has

been observed in alkaloids, quassinoids and sesquiterpene lactones. The active compounds isolated from antimalarial plants have been compiled from the review work of Saxena and others.<sup>[4]</sup> Plants which produce different antimalarial compounds, namely, alkaloids, quassinoids, sesquiterpenes, triterpenoids, flavonoids, etc. can be very important sources of antimalarial drugs. These compounds have low, moderate or high *in vitro* antiplasmodial activity, whereas some of them are inactive. They also gave a critical account of crude extracts, essential oils and active constituents with diverse chemical structure from higher plants possessing significant antimicrobial activity.

In the information obtained, also, there were many details

**Table 2: Mosquito repellent reported from northeast India**

Name of the plant	Family	Vern. name	Parts used	Methodology	References
<i>Homalomena aromatica</i> Schott.	Araceae	<i>Sugandhamantri</i> (Ass)	Rhizome	The burnt smoke of dried rhizome is used as mosquito repellent	[12]
<i>Ocimum gratissimum</i> L.	Lamiaceae	<i>Ramtulsi</i> (M., H and Beng.)	Plant	Used as mosquito repellent	[12]
<i>Elsholtzia blanda</i> Benth.	Lamiaceae	<i>Pheiri Bantulsi</i> (Ass)	Leaf	Leaf juice is used	[14], [25]
<i>Eucalyptus globules</i> Labill.	Myrtaceae	<i>Nasik</i> (M)	Leaf		[10]

(Ass) - Assamese, (H)- Hindi, (Beng)- Bengali, (M) - Manipuri, (Mi) - Mizo, (S)- Sanskrit

about the appropriate indication of each plant. For example, some plants are indicated to increase others' potency. There are also plants that are traditionally employed for specific symptoms or conditions that often accompany malaria, such as weakness, renal failure, body pain or cerebral malaria. Many plants that have been considered to lack activity against malaria due to absence of *in vitro* activity against *Plasmodium* can have other mechanisms of action. Some authors have underestimated the traditional plants used for malaria based exclusively on low activity against *Plasmodium in vitro* or in animal models. This can be a mistake of strategy or even methodology. There are many explanations for the absence of *in vitro* activity of an effective antimalarial drug. As an example, the active principle could be formed by hepatic metabolism, or as a result of transformation by gut bacteria. Other possible mechanisms of action include immunomodulation or interference with the invasion of new red blood cells by parasites, which can be species specific. Therefore, studies in human subjects, as well as the observance of ethnomedical detailed data, are required in order to exclude or confirm the activity of plants traditionally used to treat malaria.

Sixty-eight plants have been reported to act against malaria, either to kill *Plasmodium* sp. (e.g., *Al. scholaris*, *O. sanctum*) or to act as hepatic protector if used in combination with some other plant like *An. paniculata* *Co. teeta* or *Swertia chirayita*. This needs thorough screening by testing for active principles, toxicity of the extract and their pharmacological action to act against *Plasmodium* as well as hepatic protection. Therefore, it is required to carry out all possible studies on the selection basis of plants for strengthening and establishing them as a real drug before undertaking clinical trials. The plants which are commonly reported by different authors from different parts of northeast, such as *Al. scholaris*, *Co. teeta*, *Cr. occulta*, *O. sanctum*, *P. persicariaefolia*, *V. peduncularis*, should be given the priority for *in vitro* and *in vivo* studies.

There is a need to generate reliable scientific data to determine whether the plants currently used to treat malaria are actually effective. In the long term, this should help to prevent deaths due to ignorance and the misuse of plants for self-medication in the absence of advice from a qualified medical professional. Individual plants are rarely

used alone. In most cases, they are used as mixtures like *Al. scholaris* with *An. paniculata* and *Artemisia nilagarica* with *Co. teeta* and *V. peduncularis* with *An. Paniculata*, etc. It will never be easy to determine which plants are likely to be the most useful and should be examined to isolate pure active compounds. Some antimalarial plants are used for preparing baths or for inhalations (aromatic plants). It might, therefore, be useful to test the antibacterial, antiviral and anti-inflammatory activities of these groups of plants.

## CONCLUSION

The present survey has provided information about the range of species of plants used in the treatment of malaria in northeast India. (Accordingly, researchers should observe ethnomedical information of all species before deciding which kind of screening should be used in the search for antimalarial drugs.) It develops good scope for Pharmaceutics to develop new drug for malaria after combining drugs having action against *Plasmodium*, anti-inflammatory drugs as well as hepatic protector by using these traditional information and furnishing chemical analysis, pharmacological action, and *in vitro* studies. Traditional healers working in very remote parts of the region are paying much attention to treat various kinds of ailments. While using herbs for treatment, it has been observed that some are really devoted to the methodology of treatment, whereas some others concentrate on their use.

There is a need to generate reliable scientific data to determine whether the plants currently used to treat malaria are actually effective. In the long term, this should help to prevent deaths due to ignorance and the misuse of plants for self-medication in the absence of advice from a qualified medical professional. Individual plants are rarely used alone. In most cases, they are used as mixtures. It will never be easy to determine which plants are likely to be the most useful and should be examined to isolate pure active compounds. Some antimalarial plants are used for preparing baths or for inhalations (aromatic plants).

## ACKNOWLEDGMENTS

The authors acknowledge the support of Director General,

Shankar, *et al.*: Antimalarial plants of northeast India

CCRAS, New Delhi, in conducting this work. Financial assistance rendered by the Department of AYUSH, Ministry of Health and Family Welfare, Govt. of India, has also been thankfully acknowledged.

## REFERENCES

1. Symth JD. Animal Parasitology. Cambridge, England: Cambridge University Press; 1994.
2. Srisilam K, Veersham C. Antimalarials of plant origin. In: Khanum, A Khan IA editors. Role of Biotechnology in Medicinal and aromatic plants. Hyderabad, India: Ukaaz Publishers; 2003; 7:17-47.
3. Bharel S, Gulati A, Abdin MZ, Srivastava PS, Jain SK. Structure, biosynthesis and function of artemisin. *Fitoterapia* 1996;67:387-99.
4. Saxena S, Pant N, Jain DC, Bhakuni RS. Antimalarial agents from plant sources. *Curr Sci* 2003;85:9.
5. Abbiw DK. Useful plants of Ghana. London: Intermediate Technology Publications and the Royal Botanic Gardens Kew; 1990.
6. Mshana RN, Abbiw DK, Addae-Mensah I, Adjanouhoun E, Ahyi MR, Ekpere JA, *et al.* Traditional medicine and Pharmacopoeia; contribution to the revision of ethnobotanical and floristic studies in Ghana. Ghana: Science and Technology Press, CSIR; 2001.
7. Rama Shankar, Rawat MS. Medicinal plants vis a vis forest management in Arunachal Pradesh: Ayurveda and Drugs for all. 2007, Himalayan Publisher, New Delhi & Itanagar., p. 110-8.
8. Rama Shankar, Rawat MS. Ethno-midico-botanical aspect of some plants of Arunachal Pradesh. *Bull Medico Ethno Bot Res* 1995;16:83-9.
9. Rama Shankar, Rawat MS, Singh VK. Ethnomedicinal plants of Papumpare district: Ethnomedicines of the tribes of Arunachal Pradesh. In: Mibang T, editor. New Delhi & Itanagar: Himalayan publishers; 2003. p. 30-5.
10. Sharma UK. Medicinal plant of Assam. In: Singh B, Singh MP. Dehradun:2004.
11. Sinha SC. Medicinal Plants of Manipur, Mass and Sinha, Imphal: Manipur Cultural Integration Conference; 1996.
12. Lalramghinglova H. Ethno medicinal plants of Mizoram. Singh B, Singh MP. Dehradun: 2003.
13. Malhotra SC. Pharmacological Investigations of certain medicinal plants and compound formulations used in Ayurveda and Siddha. 1st ed. New Delhi: Central Council for Reseach in Ayurveda and Siddha (CCRAS); 1996. p. 409-13.
14. Anonymus. An Appraisal of Tribal folk medicines. New Delhi, India: Central Council for Research in Ayurveda and Siddha; 1999. p. 331-5.
15. Biswas K, Chopra RN. Common medicinal plants of Darjeeling and the Sikkim Himalayas. Vivek Vihar, Delhi: Periodical experts book agency; 1982.
16. Borthakur SK, Gogoi PD. Household remedies of Tai Khamptis: Ethnomedicines of the tribes of Arunachal Pradesh. Mibang T, editor. Himalayan publishers; 2003. New Delhi & Itanagar. p. 90-8.
17. Rama Shankar, Rawat MS. Medicinal plants and some folklore from East and West Siang district Arunachal Pradesh) their utilization. *Bull Medico Ethno Bot Res* 1996;17:1-7.
18. Purkayastha P, Dutta M, Nath SC. Ethno-medicinal plants from Dibru-Saikhowa biosphere reserve, Assam. *Ind J of Trad Knowl* 2007;6:477-80.
19. Issar RK. Medico-Ethno-Botanical Exploration of Sikkim Himalayas. New Delhi, India: Central Council for Research in Ayurveda and Sidha: 1991. p. 117-21.
20. Das AK, Tag H. Ethnomedicinal studies of the Khamti tribe of Arunachal Pradesh. *Indian J Tradit Knowl* 2006;5:317-22.
21. Mahanti N. Tribal ethno-botany of Mizoram, tribal Studies of India Series T 171. New Delhi, India: Inter India Publication; 1994.
22. Das AK. Some notes on the folk medicines of the Adis of Arunachal Pradesh. Ethnomedicines of the tribes of Arunachal Pradesh. In: Mibang T. Himalayan publishers; 2003, New Delhi & Itanagar p. 41-8.
23. Lalfakzuala R, Lalramghinglova H, Kayang H. Ethnobotanical usage of plants in western Mizoram. *Indian J Tradit Knowl* 2007;6:486-93.
24. Kumar S. The Medicinal Plants of North East India. Jodhpur, India: Scientific Publishers; 2002.
25. Sikdar M, Dutta U. Traditional Phytotherapy among the Nath People of Assam. *Ethno-Medicine* 2008;2:39-45.
26. Kohli YP. Ethnomedicines in the service of tani tribes. In: Mibang T, Chaudhuri SK. Ethnomedicines of the tribes of Arunachal Pradesh. Itanagar: Himalayan publishers; 2003. p. 13-7.
27. Yonggam D. Ethno Medico-Botany on the Mishing tribe of East Siang district of Arunachal Pradesh. *Arun Forest News* 2005;21:44-9.
28. Singh RK. *Tinospora cordifolia* as an adjuvant drug in the treatment of hyper-reactive malarious splenomegaly--case reports. *J Vector Borne Dis* 2005;42:36-8.
29. Karnik SR, Tathed PS, Antarkar DS, Gidse CS, Vaidya RA, Vaidya AD. Antimalarial activity and clinical safety of traditionally used *Nyctanthes arbor-tristis* Linn. *Indian J Tradit Knowl* 2008; 7(2): 330-34.
30. Wong SK, Lim YY, Abdullah NR, Nordin FJ. Assessment of antiproliferative and antiplasmodial activities of five selected Apocynaceae species. *BMC Complement Altern Med* 2011;11:3.

**How to cite this article:** Shankar R, Deb S, Sharma BK. Antimalarial plants of northeast India: An overview. *J Ayurveda Integr Med* 2012;3:10-6.

**Source of Support:** Department of AYUSH, Ministry of Health and Family Welfare, Govt. of India, **Conflict of Interest:** None declared.

## Dispatch and return notification by E-mail

The journal now sends email notification to its members on dispatch of a print issue. The notification is sent to those members who have provided their email address to the association/journal office. The email alerts you about an outdated address and return of issue due to incomplete/incorrect address.

If you wish to receive such email notification, please send your email along with the membership number and full mailing address to the editorial office by email.