

Docosahexaenoic acid content is significantly higher in *ghrita* prepared by traditional Ayurvedic method

Kalpana S. Joshi

Department of Biotechnology, Sinhgad College of Engineering, Pune, Maharashtra, India

ABSTRACT

Background: Ghee (clarified butter) also known as *ghrita*, has been utilized for thousands of years in Ayurveda. Ghee is mostly prepared by traditional method in Indian households or by direct cream method at industry level. Ayurvedic classics mention that *ghrita* made from cow milk is superior. However, there is no scientific comparison available on preparation methods and essential fatty acids content of *ghrita*. **Objective:** To investigate fatty acid composition of *ghrita* prepared by traditional/Ayurvedic method and commercial method (direct cream method). **Materials and Methods:** Fatty Acid Methyl Esters (FAME) extracted from *ghrita* samples were analysed on Gas Chromatography (GC) Shimadzu B using capillary column BPX70 (0.32 mm*60 m, ID of 0.25 mm). The fatty acids in the samples were identified by comparing peaks with the external standard 68A (Nu-Chek-Prep, Inc.USA). Significant differences between the experimental groups were assessed by analysis of variance. **Results:** Distribution of fatty acids was compared in *ghrita* samples prepared by traditional method and direct cream method which is commercially used. Saturated fatty acids were predominant in both the groups. Mono unsaturated fatty acids and poly unsaturated fatty acids were in the range of 17-18% and 3-6% respectively. DHA content was significantly higher in ghee prepared by traditional method using curd starter fermentation. **Conclusion:** The findings suggested that *ghrita* prepared by traditional ayurvedic methods contains higher amount of DHA; Omega-3 long-chain polyunsaturated fatty acids, which is a major component of retinal and brain tissues and remains important in prevention of various diseases.

Key words: Ayurveda, Docosahexaenoic acid, *ghrita*, Omega-3 fatty acids, traditional method

INTRODUCTION

Ghee or *Ghrita* is obtained by clarification of milk fats by heating and has longer shelf life at room temperature as compared with butter. Fatty acid profile of ghee is complex and shown to contain high levels of saturated fats and cholesterol. Composition varies from source of ghee such as goat milk, sheep milk, cow milk, and buffalo milk.^[1,2]

Address for Correspondence:

Dr. Kalpana S. Joshi, Department of Biotechnology, Sinhgad College of Engineering, Pune, Maharashtra, India. E-mail: joshikalpana@gmail.com

Received: 26-Sep-2013

Revised: 14-Nov-2013

Accepted: 25-Nov-2013

Ghee is mostly prepared by traditional method in Indian households or by direct cream method at industry level.^[3] The quality of ghee depends on the type and quality of milk and method of preparation.^[4,5] Ghee is fairly shelf-stable largely because of its low moisture content and possible anti-oxidative properties. Shelf life of ghee is also dependent on the method of preparation. The storing quality of desi ghee is better than that of direct cream or creamery butter ghee because of the presence of phospholipids.^[6]

The physical and chemical characteristics of the butter and ghee from cow, goat, and sheep milk are reported.^[1,7] Ghee contains 99-99.5% fats and less than 1% moisture, unsaponifiable matter and traces of charred casein, carotene, and fat-soluble vitamins.^[8] Fatty acid profile determines many of the physical, organoleptic, and nutritional characteristics of milk and dairy products.^[9] Milk fat contains over 400 individual fatty acids and their isomers. Cow milk contains large amounts of saturated fatty acids (SFA), particularly C14:0 and C16:0, and small amounts of mono unsaturated fatty acids (MUFA), poly unsaturated fatty acids (PUFA) and omega-3 fatty acids with beneficial effects on human health.^[10]

Access this article online

Quick Response Code:



Website:

www.jaim.in

DOI:

10.4103/0975-9476.131730

Many attempts have been made to alter the fatty acid composition of milk fat from dairy cows to improve the long chain poly unsaturated fatty acids (LCPUFA) content.^[11,12] The biosynthesis of LCPUFAs such as docosahexenoic acid (DHA) and arachidonic acid (AA) takes place by desaturation and elongation of precursors such as α -linolenic acid (ALA, 18:3n-3) and linoleic acid (LA, 18:2n-6), respectively. Precursors ALA and LA supplied with forage undergo bioconversion in rumen and mammary glands forming DHA and AA.^[9] Major sources of DHA include fish such as mackerel, salmon, tuna, fish oil, and deep-sea algae.^[13]

Ghee also known as “*Ghrita*” is extensively used in Ayurveda practice. Ghee is strongly recommended in daily diet and Ayurvedic therapeutics such as “*Panchakarma*” or “*Anupana*” (Vehicle) for various medicines. Traditionally ghee is made from cow milk or any other milk after churning curdled whole milk, separating the butter after fermentation, and clarifying it by heating it in pan on low flame. Alternatively, it is prepared by clarifying cream collected from raw whole milk at industrial scale. Ayurvedic classics describe eight kinds of ghee from eight different animal milk; among them ghee made from cow milk is said to be the superior.^[14] The present study was carried out to investigate fatty acid composition of “*ghrita*” prepared by traditional household/Ayurvedic method^[15] and commercial method (direct cream method).^[8]

MATERIALS AND METHODS

Preparation of “*Ghrita*” by traditional ayurvedic method

Cow milk was boiled and cooled at room temperature. About 10% curd was added as starter culture, mixed, and incubated for 8-10 hrs at room temperature. Curd formed was manually churned until butter floated on top of the buttermilk. Butter was washed 3-4 times using water to remove residual butter milk and then heated in stainless steel container till it is clarified. Clear liquid “*ghrita*” was then decanted in a glass container.

Preparation of “*Ghee*” by direct cream method

Cream is separated from whole cow milk and heated in stainless steel pan. Clear molten “*ghrita*” is decanted and stored in glass container.

Analysis of fatty acids

Modified Bligh and Dyer method was used for fatty acid analysis.^[16] Briefly, 0.1 ml of molten “*ghrita*” was taken into glass test tube, followed by addition of 10 ml of 0.6 N methanolic HCl. Samples were incubated at 80°C for 2 hrs. Samples were extracted with hexane three times. Hexane extracts were mixed together and concentrated under vacuum evaporation. Analysis of fatty acid methyl

esters (FAME) was done on GC Shimadzu B using capillary column BPX70 (0.32 mm*60 m, ID of 0.25 mm). The fatty acids in the samples were identified by comparing peaks with the external standard 68A (Nu-Chek-Prep, Inc. USA).^[17]

Statistical analysis

Data was analyzed using the Statistical Package for the Social Sciences (SPSS software version: 11, Chicago, IL). Significant differences between the experimental groups were assessed by analysis of variance. All data are expressed as mean \pm standard error of mean (SEM); P value less than 0.05 was considered to be significant.

RESULTS

Distribution of fatty acids was compared with *ghrita* samples prepared by traditional method and direct cream method, which is commercially used. Saturated fatty acids were predominant fatty acids in both the groups. Mono unsaturated fatty acids (MUFA) and poly unsaturated fatty acids (PUFA) were in the range of 17-18% and 3-6%, respectively. No significant difference was observed between the two acids [Table 1].

Distribution of selected n-6 and n-3 fatty acids was analyzed in *ghrita* samples prepared by both the methods. It was observed that there was no significant difference in precursor fatty acids such as LA and ALA contents in *ghrita* samples. DHA content was significantly higher in ghee prepared by traditional method using curd starter fermentation [Table 2].

DISCUSSION

DHA contributes to reduced risk of diseases like heart attack, cancer, insulin resistance, arthritis,^[18] and ADHD.^[19] Ghee is generally used in Ayurveda. According to Ayurveda classics *ghrita* promotes longevity and protects body from various diseases.^[20] It increases digestive fire *agni* and improves absorption and assimilation. It nourishes *dhatus*, improves memory, and has lubricating activity in joints. Most of these activities are known to be imparted

Table 1: Distribution of fatty acids in *Ghrita* prepared by traditional method and direct cream method

Fatty acid % distribution	<i>Ghrita</i> prepared by traditional method	<i>Ghrita</i> prepared by direct cream method
SFA	72.4	73.2
MUFA	18.6	17.5
n-6	5.38	6.3
n-3	3.7	3.1
n-3: n-6 ratio	0.69	0.5

SFA=Saturated fatty acids, MUFA=Mono unsaturated fatty acids

Table 2: Distribution of selected Omega-6 and Omega-3 fatty acids in Ghrita prepared by traditional method and direct cream method (n=3)

Fatty acid % distribution	Ghrita prepared by traditional method	Ghrita prepared by direct cream method
LA	5.1±0.544	6.2±1.29
AA	0.157±0.061	0.132±0.037
ALA	3.66±0.88	3.0±0.7
AA	0.218±0.096	0.169±0.95
DHA	0.083*±0.003	0.062±0.002

Values are Mean±SEM, *Value significant with $P \leq 0.05$, LA=Linoleic acid, AA=Arachidonic acid, ALA=α-linolenic acid, DHA=Docosahexaenoic acid

by anti-oxidants and essential fatty acids such as DHA. We observed consistent and reproducible increase in DHA content in ghee prepared by traditional method in controlled laboratory conditions, however, this needs to be confirmed at field level. Whether traditional method of ghee preparation protects unsaturated fatty acids from trans fat formation is unknown. Limitation of this study is that the trans fats and anti-oxidant levels of ghee samples prepared by both the methods were not measured.

We observed increase in DHA content of *ghrita* prepared by traditional fermentation method, which may be attributed to rich microbial flora used in starter curd culture. Lactic acid bacillus cultures along with added lactose and fructose are shown to promote formation of conjugated linoleic acid.^[21] Further analyses of microbial community and fatty acid profile are required to establish presence of desaturase enzyme producing bacteria responsible for conversion of precursor fatty acid ALA to DHA. DHA along with fat-soluble vitamins, anti-oxidants, and conjugated linoleic acid (CLA) could be responsible for health benefits of *ghrita*.

There are other considerations for the traditional processing of ghee. For example, mango seed kernels have been used to increase the shelf life of ghee. It has now been found that phenolics and phospholipids isolated from mango seed kernel, when added jointly to buffalo ghee, prevented peroxidation of fats and helped in extending the shelf life of ghee.^[22] Also, Ayurveda mentions about enhanced therapeutic qualities of very old cow ghee. It will be interesting to study the composition of very old cow ghee as well as physicochemical and organoleptic characteristics at various time intervals. Use of traditional ayurvedic methods to prepare *ghrita* needs to be studied for its possible role in promoting health.

CONCLUSION

The findings suggested that ghrita prepared by traditional ayurvedic method contains higher amount of DHA with beneficial effects on human health.

ACKNOWLEDGEMENT

The author would like to thank Dr. Bhushan Patwardhan for his encouragement to undertake this work. Thanks are also due to Ms. Shalini Jaiswal, M. Sc student for her technical help.

REFERENCES

- Sawaya WN, Khan P, Al-Shalhat AF. Physical and chemical characteristics of ghee and butter from goat's and sheep's milk. *Food Chem* 1984;14:227-32.
- Ramamurthy MK, Narayanan KM. Fatty acid composition of buffalo and cow milk fats by gas-liquid chromatography (GLC). *Milchwissenschaft* 1971;26:693.
- Ganguli NC, Jain MK. Ghee: Its chemistry, processing and technology. *J Dairy Sci* 1973;56:19-25.
- Rachana CR, Nath BS, Reshma AM, Armughan C. Variation in grainy texture of commercial ghee in relation to laboratory ghee and its blends. *J Int Dairy Technol* 2012;6:1-8.
- Suwarat N, Tungjaroenchai W. Characteristic of ghees obtained from different post-clarification temperatures. *Int J Biosci Biochem Bioinform* 2013;3:332-4.
- Ramamurthy MK, Narayanan KM, Bhalerao VR. Effect of phospholipids on the keeping quality of ghee. *Indian J Dairy Sci* 1968;21:62.
- Kirazci A, Javidipour I. Some chemical and microbiological properties of ghee produced in Eastern Anatolia. *Int J Dairy Technol* 2008;61:300-6.
- Srinivasan MR, Anantakrishnan CP. *Milk products of India*. ICAR, New Delhi, 1964.
- Chilliard Y, Ferlay A, Doreau M. Effect of different types of forages, animal fat or marine oils in cow's diet on milk fat secretion and composition, especially conjugated linoleic acid (CLA) and polyunsaturated fatty acids. *Livest Prod Sci* 2001;70:31-48.
- Kennelly JJ. The fatty acid composition of milk fat as influenced by feeding oilseeds. *Anim Feed Sci Tech* 1996;60:137-52.
- Collomb M, Sollberger H, Butikofer U, Sieber R, Stoll W, Schaeren W. Impact of a basal diet of hay and fodder beet supplemented with rapeseed, linseed and sunflower seed on the fatty acid composition of milk fat. *Int Dairy J* 2004;14:549-59.
- Kitessa SM, Gulati SK, Simos GC, Ashes JR, Scott TW, Fleck E, *et al*. Supplementation of grazing dairy cows with rumen-protected tuna oil enriches milk fat with n-3 fatty acid without affecting milk production or sensory characteristics. *Br J Nutr* 2004;91:271-7.
- Warude D, Joshi K, Harsulkar A. Polyunsaturated fatty acids: Biotechnology. *Crit Rev Biotechnol* 2006;26:83-93.
- Vagbhatta. In: *Ashtang Hridaya, Sutra Sthana*, 5/39. Reprinted ed. Harishastri Paradkar Vaidya., editor. New Delhi: Rashtriya Sanskrit Samsthana; 2002. p. 73-4.
- Sharangadhara, Sharangadhara Samhita, Vidyasagar PS., editors. Varanasi: Choukhambha Orientalia; 1983. p. 137-214.
- Bligh EG, Dyer WJ. A rapid method of total lipid extraction and purification. *Can J Biochem Physiol* 1959;37:911-7.
- Brenna JT, Salem N, Sinclair AJ, Cunnane SC; International Society for the Study of Fatty Acids and Lipids. Alpha linolenic acid supplementation and conversion to n-3 long-chain polyunsaturated fatty acids in humans. *Prostaglandins Leukot Essent Fatty Acids* 2009;80:85-91.
- Simopoulos AP. The importance of the omega-6/omega-3 fatty acid ratio in cardiovascular disease and other chronic diseases. *Exp Biol Med (Maywood)* 2008;233:674-88.
- Joshi KS, Lad S, Kale M, Pandit A, Mahadik S, Patwardhan B. Supplementation with flax oil and vitamin C improves the outcome of Attention Deficit Hyperactivity Disorder (ADHD). *Prostaglandins Leukot Essent Fatty Acids* 2006;74:17-21.


Joshi: DHA in ayurvedic ghrita

20. Agnivesha. In: Charaka, Dridhabala, Charaka Samhita, Sutra Sthana, Vividha Shitapitiya Adhyaya 28/45. 5th ed. Vaidya Jadavaji Trikamji Acharya., editor. Varanasi: Chaukhamba Surbharti Prakasana; 2008. p. 181.
21. Tung YL. Conjugated linoleic acid concentration as affected by lactic cultures and additives. Food Chem 2000;69:27-31.
22. Puravankara D, Boghra V, Sharma R. Effect of antioxidant principles isolated from mango (*Mangifera indica* L) seed

kernels on oxidative stability of buffalo ghee (butter-fat). J Sci Food Agric 2000;80:522-6.

How to cite this article: Joshi KS. Docosahexaenoic acid content is significantly higher in *ghrita* prepared by traditional Ayurvedic method. J Ayurveda Integr Med 2014;5:85-8.

Source of Support: Nil, **Conflict of Interest:** None declared.



Announcement

iPhone App



Download
**iPhone, iPad
application**

FREE

A free application to browse and search the journal's content is now available for iPhone/iPad. The application provides "Table of Contents" of the latest issues, which are stored on the device for future offline browsing. Internet connection is required to access the back issues and search facility. The application is Compatible with iPhone, iPod touch, and iPad and Requires iOS 3.1 or later. The application can be downloaded from <http://itunes.apple.com/us/app/medknow-journals/id458064375?ls=1&mt=8>. For suggestions and comments do write back to us.